

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

aSF69
.AIUS

United States
Department of
Agriculture ⁰²

Animal and
Plant Health
Inspection
Service

Veterinary
Services

APHIS 91-39

National Tick Surveillance Program

Calendar Year 1983

APR 28 1983

MAY 8 '85

U.S. DEPT. OF AGRICULTURE
NATL. AGRI. LIBRARY
6000 WOODBURN

National Tick Surveillance Program Calendar Year 1983

During calendar year 1983, the collection and submission of ticks from native and imported animals plus plant and animal material was 12 percent greater than in 1982. There were 10,207 collections in 1983, 9,086 in 1982, 9,381 in 1981, 7,763 in 1980, and 11,553 in 1979.

Lyme Disease

Lyme disease is a tick-borne disease of humans transmitted by *Ixodes dammini* which is characterized by a skin lesion (erythema chronicum migrans) that may be accompanied by headache, stiff neck, fever, malaise, fatigue, aching muscles and joints, and swollen lymph nodes. Weeks or months later some patients develop brain and heart infections, central and peripheral nervous disorders, and migratory musculoskeletal pain. Still later arthritis may develop and persist for several years resulting in erosion of the cartilage and bone.¹

The disease was first recognized in 1975 in children in Lyme, Connecticut. The rural setting and seasonal nature of the cases suggested possible arthropod involvement. Investigations soon incriminated a newly recognized tick species *Ixodes dammini* as one of the vectors. Interestingly *I. dammini* has in recent years been shown to vector another disease in humans. In 1976 Spielman incriminated *I. scapularis* (later recognized as the new species, *I. dammini*) in the transmission of *Babesia microti* in humans on Nantucket Island, Massachusetts.²

The etiologic agent for Lyme disease was suspected as possibly being bacterial when it was noted that when penicillin or tetracycline was given early in the illness the duration of the skin lesion was shortened and the arthritis was either prevented or attenuated. In 1982 a new spirochete was isolated from *I. dammini* in New York. In 1983 the same spirochete was isolated from the blood, skin, and cerebrospinal fluid of patients ill with Lyme disease.

In 1983 three employees of Plant Pest Quarantine, Otis Methods Development Center, Otis Air Force Base, Massachusetts, were diagnosed as having Lyme disease. The three cases were recognized in late June and early July 1983 in employees involved with testing

Gypsy Moth pheromones at Otis Air Force Base. Reportedly these clinical cases coincided with the period of greatest seasonal activity for ticks in that area.

Since Lyme disease is now known to occur in at least 14 States, Europe, and Australia, it is important to be aware of the etiology, transmission, symptoms, clinical signs, and treatment of the disease. Other endemic tick species may possibly play a significant role in the maintenance and transmission of the disease agent.

Update on Heartwater in the Caribbean

Throughout 1983 there was continued interest and concern regarding the presence of heartwater, the tick-borne disease caused by the rickettsial organism, *Cowdria ruminantium*. This disease was reported from Guadeloupe in 1980. In late 1983 official correspondence was received in Washington, D.C. from the Chief Veterinary Officer of Antigua confirming the presence of heartwater in livestock on Antigua.

Other islands in the West Indies, such as Puerto Rico, St. Kitts, and Martinique, are especially concerned since they also presently have established populations of *Amblyomma variegatum*, a primary vector of heartwater. In 1983 there were 39 laboratory confirmed collections of *A. variegatum* from Puerto Rico. It is strongly emphasized, however, that there were no reports of heartwater from Puerto Rico in 1983 or earlier years.

There is a potential threat for the introduction of heartwater into the continental United States since one of the experimental vectors, *Amblyomma maculatum*, is endemic in 10 States. These are primarily Gulf of Mexico or Atlantic Ocean coastal States. Increased illicit drug and animal traffic, along with the influx of illegal aliens, increases the threat of introducing *A. variegatum* and heartwater in coastal States such as Florida. It is important that Federal and State regulatory officials improve the surveillance program to rapidly detect and eliminate heartwater should it be introduced onto the mainland of the United States.

African Swine Fever

In light of the recently demonstrated potential for the soft tick, *Ornithodoros puertoricensis*, to transmit swine fever (ASF) virus, a great deal of interest has centered around this tick.³ In July 1983 a team of USDA, Agricultural Research Service (ARS), and Animal and Plant

¹Steer, Allen C. et al. 1983. The Spirochetal Etiology of Lyme Disease. New England J. Med., Vol. 308, No. 13:733-742.

²Spielman, Andrew. 1976. Human Babesiosis on Nantucket Island: Transmission By Nymphal *Ixodes* Ticks. Am. Jour. Trop. Med. Hyg. Vol. 25, No. 6: 784-787.

³Butler JF, and Gibbs EPJ. 1983. Distribution of Potential Soft Tick Vectors of African Swine Fever in the Caribbean Region. Prev Vet Med: In press.

Health Inspection Service (APHIS) entomologists conducted a survey for *O. puertoricensis* in the Dominican Republic. Surveying for this burrow-dwelling tick was greatly enhanced due to the use of a vacuum sampling device developed by researchers of the University of Florida at Gainesville. Serological evaluation, using an agar gel double-diffusion test to determine host blood present in the ticks, was performed at the Pathobiology Laboratory of the National Veterinary Services Laboratories, (NVSL), Ames, Iowa. No ticks were shown to have fed upon hogs, although 20 percent of those collected had fed on rats and mongooses. ASF virus has not been isolated from samples taken in the Dominican Republic in 1983. The eradication effort to curb this highly contagious viral disease appears to be progressing satisfactorily.

Status of ARS Cattle Fever Tick Research

In October 1983 the new USDA, ARS Cattle Fever Tick Research Laboratory was opened at Moore Field near Mission, Texas. The quarantined facility is situated within double security fences and occupies about 103 acres of land. The new laboratory will provide opportunities for field-type experiments that were not possible at the old laboratory site at Falcon Heights, Texas. The laboratory complex includes a general laboratory and office building, a separate acaricide laboratory building, barns with stalls to accommodate up to 64 animals, a covered experimental dipping vat and spray application area, a program dipping vat, and a utility building. Future research efforts at Mission will include studies to evaluate the susceptibility to acaricides of fever ticks collected from tick outbreaks within the Tick Eradication Quarantine Area, which separates Texas from Mexico. Other areas of research shall include the following:

- (1) acaricide testing as warranted by special problems or the need to test selected new chemicals,
- (2) studying the role of white-tailed deer and exotic game in the ecology of fever ticks,
- (3) expanded efforts to perfect the sterile hybrid *Boophilus* method of tick eradication, and
- (4) a variety of other projects related to the ecology and eradication of *Boophilus* ticks. Dr. Ronald B. Davey is the scientist-in-charge of the laboratory.

During 1983 Dr. Glen I. Garris and his staff of the ARS Tropical Tick Research Laboratory at Mayaguez, Puerto Rico, completed field evaluations of fenvalerate, permethrin, and amitraz. None of these three acaricides is

as effective against adult *B. microplus* as organophosphates such as coumaphos or crotoxyphos, but they are very efficacious against immature ticks. Spray treatments at the manufacturer's recommended concentrations provide almost 100 percent protection for 4-7 days' posttreatment against reinfestation of treated cattle by larvae. Tick research in Puerto Rico also includes the second year of a 2-year study of *B. microplus* ecology. When this investigation is completed in December 1984, it will confirm the survival times of larvae in both high and low rainfall areas on the island.

Dr. Garris has begun a cooperative study of *Amblyomma variegatum* with French scientists on Guadeloupe. They will evaluate the susceptibility of *A. variegatum* to acaricides and determine factors that influence its distribution in the Caribbean. Studies will also be conducted to determine the survival rate of free-living stages of the tropical bont tick.

Experimental Transmission of Anaplasmosis by Males of *Dermacentor albipictus* and *D. occidentalis*

Agriculture Research Service scientists recently unequivocally incriminated the males of *Dermacentor albipictus* and *D. occidentalis* as intrastadial, biological vectors of *Anaplasma marginale* under experimental conditions.⁴ This is apparently the first such record for the males of a one-host tick species such as *D. albipictus*. An earlier preliminary study by Stiller had suggested that the males of the one-host tick, *D. albipictus*, could serve as competent intrastadial, biological vectors of the causative agent of anaplasmosis.

The confirmation of the role of one-host males to act as intrastadial, biological vectors of *A. marginale* has potential significance because if such males transfer to more than one host animal in the field, they could acquire and transmit the parasite in the absence of infection in the host animal on which they fed as immatures. This could increase the vector potential of one-host ticks, perhaps including important vector species such as *Boophilus microplus* and *B. annulatus*.

The males of the three-host species, *D. occidentalis*, are especially efficient vectors of *A. marginale* as demonstrated in the experiment where as few as three *D. occidentalis* males transmitted the causative agent. This may be especially important in that it has been often assumed that *A. marginale* is transovarially

⁴Stiller, D.; Johnson, L. W.; and Kuttler, K. L. 1983. Experimental Transmission of *Anaplasma marginale* Theiler by Males of *Dermacentor albipictus* (Packard) and *Dermacentor occidentalis* Marx (Acari: Ixodidae). Proc. US Animal Health Assoc., pp. 59-65.

transmitted by the female of *D. occidentalis* although this assumption is based on a single study conducted in the mid-1930's. This early study did not address the possibility of accidental fly transmission of *A. marginale* and subsequent studies were unable to repeat the transovarial transmission of *A. marginale* by *D. occidentalis*. For this reason, and because the common hosts of the immature stages of this tick are not known to harbor *A. marginale*, the males of *D. occidentalis* could represent an important means by which *A. marginale* enters the three-host cycle of this vector species.

The true significance of male tick vector competence in the epizootiology of anaplasmosis cannot be accurately assessed without additional information on the frequency of interhost transfer by males in the field. This important aspect should receive additional research to fully determine the role male ticks may play.

***Amblyomma variegatum* Eradication in Puerto Rico**

The tropical bont tick continues to be a problem in Puerto Rico. There were 38 laboratory-confirmed collections in 1983 as compared to 50 collections confirmed in 1982. The areas known to have infested livestock in 1983 include the municipalities of Cabo Rojo and Ponce and the island of Vieques. These infested areas are widely separated with Cabo Rojo on the western coast and Ponce approximately on the south central coastal area of the main island of Puerto Rico. Vieques is an island some 10 miles off the eastern coast of Puerto Rico.

***Boophilus microplus* in Puerto Rico**

Eradication efforts continued in Puerto Rico against the widespread infestation of *B. microplus*. The National Veterinary Services Laboratory, Ames, Iowa, confirmed 4,295 collections of *B. microplus* in 1983 as compared to 2,208 collections in 1982. Over 98 percent of the collections were taken from cattle with less than 2 percent of the hosts being horses and goats.

One serious problem encountered in 1983 was the reinfestation of herds recently freed of ticks and released from quarantine. Federal and Commonwealth tick eradication personnel continue to implement changes in the program to correct problems as they are detected.

***Boophilus* Tick Eradication in Texas**

Compared to recent years, the tick eradication activities for 1983 were successful and relatively uneventful. There were no major outbreaks of *Boophilus* ticks in either the Tick Eradication Quarantine Area (TEQA) or

the Free Area (FA). Only two introductions involving 543 cattle and 5 horses occurred in the FA. These introductions were quickly confined and eradicated. Several introductions occurred in the TEQA which borders on the Rio Grande. These involved small numbers of livestock and were also rapidly contained and eliminated.

Despite a reduction-in-force of 11 inspectors in the spring of 1983, the fever tick problem was minimal compared to recent years. This was primarily due to the hard work and the dedication of inspectors and the cooperation of industry people. Because of relatively few quarantines in the FA, inspectors were able to concentrate on river patrols to stop the introduction of ticks at this point.

The weather conditions in 1983 were extremely dry. Most range lands had little or no forage for livestock and some stock ponds were also dry. The Rio Grande was lower than normal allowing for easier crossing of stray Mexican livestock.

Smuggling of horses from Mexico continued to be a problem in 1983. Tick eradication personnel apprehended 19 head of horses involving 18 illegal introductions. The Tick Eradication Force was responsible for the apprehension of 73 stray cattle and 109 stray horses representing 56 separate incidents of which six involved *Boophilus*-infested animals.

Legislation has been passed giving certain USDA employees the authority to carry firearms. Firearm safety and training was initiated for all tick force employees in December 1983 at Glynco, Georgia.

Ticks and Zoological Compounds

In recent years there has been a trend to develop large, commercial zoological compounds throughout the United States. Generally, these are large, double-fenced compounds where a variety of compatible wild animals roam freely. The general theme is to develop habitats suggestive of Africa where the animals usually originate. For a fee, tourists or other interested persons may tour the facility in their own automobiles or in vehicles provided by the management. This allows them to observe the wildlife in their "native" habitat.

However, such a compound poses some problems regarding the possible introduction of exotic ticks on animals from Africa. Some species of animals such as rhinoceroses, elephants, hippopotamuses, and the big cats are not presently required to be held in postentry quarantine.


Nevertheless, these animals may serve as suitable hosts for the immature and mature stages of economically important ticks. For example, in the past rhinoceroses have entered the United States and were subsequently found infested with live *Amblyomma hebraeum*. In 1960, live *Rhipicephalus evertsi* ticks were found on elands, zebras, and nilghai in "Africa USA" at Boca Raton, Florida. An intensive eradication effort was required to eradicate this tick. Shortly thereafter, *R. evertsi* and *R. pulchellus* were found at "Busch Gardens," Florida, and at the "Catskill Game Farm" in up-state New York.

Exotic ticks are also commonly found on animals in quarantine at the USDA New York Animal Import Center, Newburgh, New York. Precautionary dipping is routinely carried out on such animals held in quarantine. The real threat exists with those animals which may enter the USA without having to be held in quarantine or having to be given a precautionary dipping. Not only are many exotic ticks serious blood-sucking parasites of domestic livestock, but they also transmit diseases such as anaplasmosis, babesiosis, theileriosis, heartwater, Nairobi sheep disease, and spirochetosis.


Procedures for Collecting and Submitting Ticks

Veterinary Medical Officers and Animal Health Technicians should carefully review V.S. Memorandum 558.1, September 8, 1978, for information on the collection and submission of ticks. Any personnel not having this memorandum may obtain a copy from their Veterinarian-In-Charge. Cooperating State animal health regulatory personnel may also obtain a copy of this memorandum.

Tick Collections From All Hosts CY 1983



Tick Collections From Animals and Products Offered For Entry CY 1983



REPORT OF TICKS COLLECTED													PERIOD		HOST													
													Calendar Year 1983		Animals and Products Offered for Entry													
STATE	TOTAL	AMBL.YOMMA AMERICANUM	AMBL.YOMMA CAJEUENSE	AMBL.YOMMA ILLIATIBEL	AMBL.YOMMA MACGILLIUM	AMBL.YOMMA SPP.	ARGAS HEFESICUS	BOOPHILLUS ANNULATUS	BOOPHILLUS MICROELLUS	BOOPHILLUS SPP.	DERMACENTOR ALDOPICINUS	DERMACENTOR AMERICANUS	DERMACENTOR V. B. VAUGHANI	DERMACENTOR MICROLINEATUS	DERMACENTOR TILLENI	DERMACENTOR OCCIDENTALIS	DERMACENTOR VERTICILLUS	DERMACENTOR SPP.	HEMAPHYSALIS PARVICTATUS	IXODES SCOPULI	IXODES PACIFICUS	IXODES SCAPULARIS	IXODES SPP.	ORNITHODOROS SPP.	OTIOBIUS MEGNINI	RHIPICEPHALUS SENSIENSIS	RHIPICEPHALUS SPP.	RHIPICEPHALUS MISC. SPECIES
TOTAL	148	10	2	2	1		24	19	24	25			3			5	12	2			1	1		12	1	3	1	
ARIZONA																												
Douglas	10									10B0																		
Tucson	5									1D							4BH ^d											
CALIFORNIA																												
Lawdale	1																											1H0 ^{aa}
FLORIDA																												
Miami Airport	1																									1B0		
MARYLAND																												
Balt/Wash Airport	1																	1DW										
NEW MEXICO																												
Columbus	4									2B0		1B0					1B0 ^k											
NEW YORK																												
JFK Airport	2																					1PM ^{ip}					1PM ^{rc}	
Rock Tavern	2																										1Z ^{re}	1Z ^{rm}
TEXAS																												
Del Rio	14	1B0					3B0	3B0	5B0**								1B0**				1B0							
Eagle Pass	41			1B0			19B0	2B0	7B0**	6B0		2B0			2B0	1B0**	1B0**											
El Paso	19									5B0							2B0**						12B0					
Laredo	47	9B0	2B0	1B0	1B0**	2B0	1D	12B0	12B0**	1B0					2B0	1H0	2B0**											
Presidio	1						1H0										1B0 ^k											

ANH Form 5-55
MAY 1971

REPLACES ADE FORM 5-55, DEC 1964, WHICH IS OBSOLETE


USDA, APHIS, VS
NVSL, AMES, IOWA

Key to Host
 BG Baggage
 BO Bovine
 BH Bighorn Sheep
 D Deer
 DW Dead Wood-wooded
 area of airport
 HO Horse
 FM Plant Material
 Z Zebra

Key to Species
 aa *Haemaphysalis longicornis*
 (= *H. bispinosa*)
 d *Dermacentor hunteri*
 k *Dermacentor parumapertus*
 ip *Ixodes pilosus*

rc *Rhipicephalus compositus*
 re *Rhipicephalus evertsi evertsi*
 rm *Rhipicephalus evertsi mimeticus*
 ** Nymphal stage
 Bracketed figures refer to Laredo

Tick Collections From Dogs CY 1983



REPORT OF TICKS COLLECTED

PERIOD

HOST

Calendar Year 1983

Dogs

STATE	TOTAL	PERIOD														HOST													
		AMBL.YOMMA AMERICANUM	AMBL.YOMMA CAJURENSE	AMBL.YOMMA INFLATATA	AMBL.YOMMA MACULATA	AMBL.YOMMA SPP.	ARGAS PERICLUS	BOOPHILLUS ANNULATUS	BOOPHILLUS MICROPLUS	BOOPHILLUS SPP.	DERMACENTOR ALBERTUS	DERMACENTOR SYMPLECTICUS	DERMACENTOR VENTRILOQUIUS	DERMACENTOR ALBERTUS	DERMACENTOR OCCIDENTALIS	DERMACENTOR VARIABILIS	DERMACENTOR SPP.	HAEMAPHYSALIS LEIDYI	HAEMAPHYSALIS SPP.	IACOPS COCKLEI	IACOPS SPP.	IACOPS SEPILLARIS	IACOPS SPP.	ORNITHODOROS SPP.	OTLOBUS MEXICANI	RHINCEPHALUS SPP.	RHINCEPHALUS SPP.	RHINCEPHALUS SPP.	MISCELLANEOUS SPP.
TOTAL	440	16	2	12	2					1				166								14	1		3	223			
Alabama	4													2													2		
Alaska																													
Arizona																													
Arkansas																													
California																													
Colorado	8									1															1	6			
Connecticut																													
Delaware																													
Florida	13	2		1										4								4				2			
Georgia	2																					1				1			
Hawaii																													
Idaho	3													1													2		
Illinois																													
Indiana																													
Iowa																													
Kansas																													
Kentucky																													
Louisiana																													
Maine																													
Maryland																													
Massachusetts																													
Michigan																													
Minnesota																													
Mississippi	2													1												1			
Missouri																													
Montana																													
Nebraska																													
Nevada																													
New Hampshire																													
New Jersey																													
New Mexico																													
New York	1	1																											
North Carolina	7	1												1								1				4			
North Dakota	2													2															
Ohio																													
Oklahoma																													
Oregon																													
Pennsylvania																													
Rhode Island																													
South Carolina	12	1												2								4	1 ^a			4			
South Dakota																													
Tennessee	1													1															
Texas	377	11	2	11	2**									151								4			2	194			
Utah																													
Vermont																													
Virginia	1													1															
Washington																													
West Virginia																													
Wisconsin																													
Wyoming																													
Puerto Rico	7																										7		
Virgin Islands																													

AMH FORM 5-55 MAY 1971


REPLACES AGR FORM 5-55, DEC 1964, WHICH IS OBSOLETE

Key to Species
a *Ixodes affinis*


** Nymphal stage

USDA, APHIS, VS
NVSL, AMES, IOWA


Tick Collections From Horses and Mules CY 1983



Tick Collections From Native Wildlife CY 1983



Tick Collections From Zoo Animals And Miscellaneous Hosts CY 1983



REPORT OF TICKS COLLECTED														PERIOD Calendar Year 1983		HOST Zoo Animals and Miscellaneous Hosts														
STATE	TOTAL	AMBL.YOMMA AMERICANUM	AMBL.YOMMA CA. TENNENSE	AMBL.YOMMA MEXICA DDR	AMBL.YOMMA MEXICOLA LM	AMBL.YOMMA SPP.	ARGAS PERGULUS	BODIPHILLUS ANNULATUS	BODIPHILLUS BUDROPULUS	BODIPHILLUS SPP.	DERMACENTOR ALBULECTUS	DERMACENTOR SIMPSONI	DERMACENTOR XENUS S & D	DERMACENTOR ALUCRUMMENTUS	DERMACENTOR VITENS	DERMACENTOR ZOOEPHALUS	DERMACENTOR VARIABLE	DERMACENTOR SPP.	HAEMAPHYSALIS LEPIDI	IXODES PALLETIUS	IXODES COCKEI	IXODES PACIFICUS	IXODES SCAPULARIS	IXODES SPP.	ORNITHODOROS SPP.	OTOBULUS MEXICANUS	RHIPICEPHALUS SANGUINEUS	RHIPICEPHALUS SPP.	MISCELLANEOUS SPP.	
TOTAL	116	13	2	2	5	5			33						5		30			1		4				2	12	1	1	
Alabama																														
Alaska																														
Arizona																														
Arkansas																														
California																														
Colorado																														
Connecticut																														
Delaware																														
Florida	8	20 1H			1H												1HM					1Q					1HS 1X			
Georgia	2				1Q																	1Q								
Hawaii																														
Idaho																														
Illinois																														
Indiana																														
Iowa																														
Kansas																														
Kentucky	2																1H 1G													
Louisiana																														
Maine																														
Maryland																														
Massachusetts																														
Michigan																														
Minnesota																														
Mississippi	2																										1X	1WH ^{FS}		
Missouri	1	1Q																												
Montana																														
Nebraska																														
Nevada																														
New Hampshire																														
New Jersey																														
New Mexico																														
New York																														
North Carolina	2	1C															1G													
North Dakota																														
Ohio																														
Oklahoma																														
Oregon																														
Pennsylvania																														
Rhode Island																														
South Carolina	3																2H						1S							
South Dakota																														
Tennessee	1																1H													
Texas	57	6H 1X	2H	2H	1C 1H	1C ¹¹ 1Q ^{**}											5C 7Y			1Q		1C			1C 1Q	3C 2Q		1PH ^{ar}		
Utah		1DK			1G	3H ^{**}											9H 1Q 1G										1H 3X			
Vermont																														
Virginia																														
Washington																														
West Virginia																														
Wisconsin																														
Wyoming																														
Puerto Rico	38								20G 13Q						5Q															
Virgin Islands																														

ANN FORM 5-55
MAY 1971

REPLACES ADE FORM 5-55, DEC 1964, WHICH IS OBSOLETE

Key to Host
 C Cat HS Horse Stall WH Wart Hog (trophy hide, origin
 DK Donkey PH Chicken House South Africa)
 G Goat Q Host not given X Inanimate object
 H Human S Sheep Y Swine
 HM Dwelling

Key to Species
 ar Argas radiatus
 n Amblyomma inornatum
 rs Rhipicephalus simul
 ** Nymphal stage

Brackets enclose Texas collections
 USDA, APHIS, VS
 NVSL, AMES, IOWA

